



## User-Friendly Features for Better Monitoring

- No coding technology and large LCD display offers easier glucose monitoring
- HI and LO messages remind patients to take necessary action
- Five customizable test reminders remind user to test at specific times



- Meal markers included to monitor results before and after meals
- Quick 5 second testing with tiny 0.8  $\mu$ L blood sample and under dosing detection
- Alternative site testing options for less pain



# On-Call<sup>®</sup> Vivid Pal

## Blood Glucose Monitoring System

Compact Design with Reliable Results for Better Care

No Coding Required



ACON Laboratories, Inc., 10125 Mesa Rim Road, San Diego, CA 92121, U.S.A.  
 Tel: 1-858-875-8000 Fax: 1-858-200-0729 E-mail: [info@aconlabs.com](mailto:info@aconlabs.com)  
 Please visit our website for details: [www.acondiabetescare.com](http://www.acondiabetescare.com)



1150710802

© 2014 ACON Laboratories, Inc.

For self testing and professional use.



## Why is SMBG important for diabetics?

Self-monitoring of blood glucose (SMBG) enables better blood glucose control, which decreases the long-term risks of diabetic complications. It allows patients to track their blood glucose levels anytime, which helps prevent the immediate and potentially serious consequences of very high or very low blood glucose levels. Self-monitoring your blood glucose levels is one of the best ways to examine how well a diabetes treatment plan is working for you.

## How often do you need to test?


SMBG Frequency Treatment Regimen	Blood Glucose Level	
	At Target	Above Target
Multiple daily injections or insulin pump	> 3-4 times/day	> 3-4 times/day
Oral agents or once-daily insulin	> 1 time/day + 1 profile*/week	> 2 times/day
Oral agents plus once-daily insulin	> 1 time/day + Frequent profiles	> 2 times/day
Non-pharmacologic therapy	> 1 profile/week	> 1 profile/week

\* A collection of pre- and post-meal glucose test results over a 7-day period.

Ref.: The American Journal of Medicine (2005) Vol 118 (Suppl 9A), 1S-6S

## What is your target blood glucose level?

The table below shows tight diabetes control targets, which are the normal target levels for nondiabetics that have been recommended by the ADA. Aiming to control the glucose levels as close as possible to the numbers below will help the diabetic to prevent complications. It is important to set your individual goals with your health care provider.



Please consult your health care provider for your blood glucose target ranges.

Ref.:Diagonosis and Classification of Diabetes Mellitus, ADA DIABETES CARE, 2011 (01 2011 ) VOL34, SUPPLEMENT 1

## Quick and Reliable Results with Advanced Technology

- No coding required
- GOD enzyme to avoid the interference from maltose, galactose, etc
- Accurate results in just 5 seconds
- Permits re-dosing of 2<sup>nd</sup> blood sample within 3 seconds when insufficient sample is added

## Convenient Operation with Compact and Stylish Design

- Large display size for easy reading of test results
- Small size makes it easy to carry and convenient for testing anytime, anywhere



- Easy, two-step, fast and accurate testing



Insert a test strip to turn on the meter



Apply a tiny 0.8 µL blood sample and obtain result in just 5 seconds